



ROSULfan™ A

Ammonium Lauryl Sulfate

Local. Global. Integrated.

Description

- an alternative to SLS and SLES
- milder effect on the skin compared to the basic anionic surfactants
- the ability to produce dense and stable foam

Application

- shampoos
- body wash products
- shower gels
- liquid soaps
- conditioners



ROSULfan™ A

Ammonium Lauryl Sulfate

Chemical name	Sulfuric acid, mono-C12-14 -alkyl esters, ammonium salts	
INCI name	Ammonium Lauryl Sulfate	
CAS number	90583-11-2	
Function	Base surfactant, foaming agent	
Technical requirements	Appearance at temperature 30°C	clear viscous liquid
	Klett colour, Klett value	max. 30
	pH of 20% solution	4.5 ÷ 6.0
	Active substance, % (m/m)	26.0 ÷ 28.0
	Unsulphated substance, % (m/m)	max. 0.6
	Ammonium sulphate (VI), % (m/m)	max. 1
General data	Density, g/mL	approx. 1.0
	Preservative	0.3% benzoic acid
	Molecular weight	approx. 294

Shampoo for damaged and fragile hair

Phase	INCI name	Brand name	Concentration [%]	Function
A	Aqua		46.62	solvent
A	Xanthan Gum		0.75	viscosity modifier
A	Glycerin		2.00	moisturising agent
A	Microcrystalline Cellulose		0.50	viscosity modifier
A	Aqua		13.00	solvent
A	Citric Acid		0.20	pH modifier
A	Polyquaternium-10		0.03	conditioning agent
B	PEG-7 Glyceryl Cocoate	ROKAcet KO300G	2.00	re-oiling agent
B	Ammonium Lauryl Sulfate	ROSULfan A	10.00	surfactant
B	Sodium Lauroyl Sarcosinate	ROKAtend LS	20.00	surfactant
B	Cocamidopropyl Betaine	ROKAmina K30	3.50	surfactant
C	Parfum		0.40	fragrance
C	Ehylhexylglycerine, Phenoxyethanol		1.00	preservative

Appearance	visual method	viscosus milky gel
pH		5.0 - 7.0
Viscosity [cP]	Brookfield LV, spindle: 34, speed: 2,5 RPM, T:25°C	9000 - 11000
Stability	1 month in 5°C, 20°C, 40°C,	confirmed

Procedure:

1. In a main vessel combine ingredients from the phase A. Add Xanthan Gum to Glycerin - mix until homogenous solution is obtained. Add warm water (50-55°C) and Microcrystalline Cellulose. Mix until homogenous solution is obtained. Homogenise for 2-3 minutes.
2. Combine ingredients from the phase B. During mixing add Citric Acid and Polyquaternium-10 to warm water (50-60°C). Mix until homogenous solution is obtained. Add the rest of the phase B components. Mix until uniform.
3. Add phase B to phase A. Mix until homogenous solution is obtained. Cool the batch down to 30°C.
4. When the batch temperature is 30°C, add parfum and preservative. Mix until uniform.



PCC Exol SA

Sienkiewicza 4

56-120 Brzeg Dolny, Poland

products@pcc.eu

Please visit our capital group business platform:

www.products.pcc.eu



September 2026

The information in the catalogue is believed to be accurate and compiled to the best of our knowledge; however, it should be considered as introductory only. Detailed information about our products is available in TDS and MSDS.

The suggestions for product applications are based on our best knowledge.

The responsibility for the use of products in conformity or otherwise with the suggested application, and for determining product suitability for the user's own purposes rests with the user.

All copyright and trademark rights, as well as other intellectual and industrial property rights and the resulting rights to use this publication and its contents have been transferred to PCC Rokita SA or PCC EXOL SA or its licensors. All rights reserved.

Users/readers are not entitled to reproduce this publication in whole or in part, nor are they entitled to reproduce it (excluding reproduction for personal use) or to transfer it to third parties.

Permission to reproduce it for personal use does not apply to data used in other publications, electronic information systems, or other media publications. PCC Rokita SA and PCC EXOL SA shall not be responsible for data published by users.